COMMONWEALTH OF KENTUCKY DEPARTMENT OF MINES & MINERALS

THIRD DRAFT OF

RULES GOVERNING MINE EMERGENCY TECHNICIAN CONTEST

OCTOBER 9, 2002

KENTUCKY DEPARTMENT OF MINES AND MINERALS MINE EMERGENCY TECHNICIAN CONTEST

Table of Contents	Page I
Contact person for questions regarding third draft of MET contest policy, rules and procedures	Page II
Policies, rules and procedures	Page III & IV
General Rules	Page 1
Contest Judging	Page 5
Written Examination	Page 6
General Rule Discounts	Page 6
General Rule 1 Clarification	Page 7
General Rule 2 Clarification	Page 7
Timing Rules	Page 8
Recommended Functions and Critical Skills	Page 8
Skill Evaluation Station	Page 8
Summary of Skills	Page 9
Individual Skills	Page 10 thru 39

This first draft of policies, rules and procedures for the Mine Emergency Technician (MET) contest was developed at the Harlan office of the Kentucky Department of Mines and Minerals by Wayne Collett, Ronnie McDowell and Jim Owens, on February 11, 1999. A second draft was completed and dated March 15, 1999.

For questions regarding this third draft, dated October 9, 2002, please contact Wayne Collett at the Harlan office of the KDMM at 606-573-1260.

EVERY

KENTUCKY

COAL

MINER

COUNTS

KENTUCKY DEPARTMENT OF MINES AND MINERALS (KDMM) MINE EMERGENCY TECHNICIAN (MET) CONTEST POLICIES AND PROCEDURES

- I. Objectives for Planning and Management of MET Contests
 A. Contest Objectives
 - (1) The MET Contest will serve as an effective instrument for training and improving the skills required of participants in an emergency response situation.
 - (2) The MET Contest will provide a forum for building relationship and understanding in the area of emergency preparedness and response among labor, management, and regulatory agencies.
 - B. MET Contest Management Structure

A Chairman shall be appointed by the Commissioner of Mines and Minerals and shall be responsible for organizing and managing the MET Contests.

- (1) A contest committee, consisting of six (6) members will also be appointed by the Commissioner, to serve with the Chairman in organizing and managing the MET contests; developing contest rules.
- (2) The contest committee will consist of three (3) employees from the Kentucky Department of Mines and Minerals and three (3) active employees from the coal industry.
- (3) The Committee shall meet at least once annually. Any member unable to attend scheduled meetings should notify the Chairman; those who fail to make notification and who miss two consecutive meetings may be removed from the Committee.
- II. Qualifications and Selection of Contest Judges

A. CHIEF JUDGE AND ASSISTANT CHIEF JUDGE(S):

- (1) Will be appointed by the Chairman.
- (2) Will be a Kentucky Department of Mines and Minerals employee possessing a valid MET instructor certificate from the Kentucky Department of Mines and Minerals.
- (3) Contact with MET team participants will be limited to rules interpretation only. This does not preclude the Chief Judge or Assistant Chief Judge(s), from conducting initial M.E.T. training

or annual refresher training for MET provided that such training does not include specific reference to MET contest rules or work.

B. JUDGES

- (1) M.E.T. Contest Judges will hold a valid MET certificate from the Kentucky Department of Mines and Minerals, or a valid EMT certificate from Kentucky Cabinet for Human Resources.
- (2) The Kentucky Department of Mines and Minerals will verify the qualifications of each potential contest judge.
- (3) The Chief Judge will select judges (including Chairman of each contest judging team) from the list of qualified candidates. The Kentucky Department of Mines and Minerals will approve the selection of judges.
- (4) The Chairman of each contest judging team will be a Kentucky Department of Mines and Minerals employee, unless otherwise specified by the Chairman of the Contest Committee.
- (5) At least once each year, prior to the MET Contest season, the Chairman will meet with all MET judges to explain all rules, procedures, changes and interpretations.
- (6) Selected judges will be trained by the KDMM prior to judging in the MET Contests. This training will include current rules interpretation and conduct.

C. CONTEST APPEALS COMMITTEE

(1) The Chief Judge will appoint the Contest Appeals Committee. Selections will be made based on MET experience and contest work.

D. CONDUCT OF JUDGES AND OTHER CONTEST OFFICIALS

- (1) Each judge/contest official will maintain the confidentiality of the problem and scoring. Judges will discuss discounts and team performance with responsible contest officials only.
- (2) Professional standards will be maintained at all times by judges and other contest officials.
- (3) Judges and contest officials will be prompt in reporting for their assigned duties, display courtesy, and will be properly dressed for their job assignment.

KENTUCKY DEPARTMENT OF MINES AND MINERALS MINE EMERGENCY TECHNICIAN CONTEST

GENERAL RULES

A MET Team shall consist of two people who are currently certified as a Mine Emergency Technician (MET) and currently certified in Cardio Pulmonary Resuscitation (CPR). An additional person may be used as a patient, alternate, or substitute. If a patient is not provided by the participating teams, one will be provided by the Chief Judge.

Mine Emergency Technician (MET) teams can enter MET contests by filing a pre-registration form and submitting the entry fee payment to the contest sponsor.

During registration at the MET contest, participating teams shall present proof of current MET and CPR certification. The teams shall also identify members who will be serving as Primary MET, Secondary MET and patient. (Patients will be provided by the Chief Judge if necessary). Changes in these assignments cannot be made after registration, unless approved by the Chief Judge. Teams will also draw numbers during registration to determine order of performance.

Teams will be required to perform problems while utilizing infection control skills and equipment. After problem completion, teams will be required to remove all used and unused material from the field, in accordance with infection control techniques and proper equipment storage. (Note: trash bags can be used in lieu of biohazard bags during the contest).

After registration, participating teams will be placed under guard at a location designated by the Chief Judge. They will remain under guard until time to work the problem.

Any team member receiving unauthorized information concerning a contest problem while under guard or traveling to their work station shall be disqualified by the Chief Judge and Final Appeals Committee.

Each participating team, once notified will have 20 minutes for reviewing the Score Cards and to prepare any protest. All protests shall be in writing and shall state the discount in question, the score card involved, and their reference proof (either the specific page in the Brady Basic First Response Text and/or the rule book) to support their protest. All protests will be considered by the final Appeals Committee. A decision by the Final Appeals Committee is binding and final. Protest sheets will be furnished to the teams by the Judges for the recording rules infractions or discounts assessed to teams.

MET teams will be judged on: recommended functions and skills, the written examination and actual work time for problem completion.

Participants will be required to furnish their own First Response Kit and the following listed articles from the MET/EMT Station. Listed below is the equipment required.

FIRST RESPONSE KIT MINIMUM EQUIPMENT LIST

4	Adhesive Tape, 1" x 10 yds.
1	Splint - Arm (full adult)
1	Splint - Leg (full adult)
1	Airways - (Oropharyngeal - 1 kit adult sizes)
1	Adult Bag Valve Mask – Assembled as much as possible)
1	Blood Pressure Cuff (Sphygmomanometer – assembled)
1	Stethoscope
1	Burn Sheet, Sterile 56" x 90" minimum
2	Cold Packs
10	Disinfectant Hand Cleaner (packets or equivalent)
1 each	Small, Medium and Large Extrication Collars or (2) Adult Adjustable Collars
2	Eye Shields
3	Infection Control Kits (which shall include, but not limited to):
1	Eye or Face shield, transparent, rigid material
1	Protective gown or coveralls, disposable
1	Protective foot covers, disposable
1	Red plastic bag (biohazard) for disposal
1	Surgeon cap, disposable
1	Surgeon face mask, disposable
1	Latex/vinyl examination gloves (1 pair per kit)
4 pairs	Latex/vinyl examination gloves (in addition to kits)
1	Nasopharyngeal Airways (1 kit – with non-petroleum gel lubricant)
10	Oval Eye Pads
1	Pocket Mask (with one-way- valve)
8	Roller Bandage Gauze, 4" x 5 yards
1	Scissors, EMT Utility

2	Space Blankets (disposable, weatherproof)
24	Sterile Gauze, 4" x 4"
1	Adult Suction Unit
4	Trauma Dressings, Sterile, Large
12	Triangular Bandages, Large
1	Potable Water, 1000 ML
5	Patient Care Report Forms and Pen or Pencil

Revised 7-17-01

The Mine Emergency Technician Team will not be allowed to bring any materials other than those stated above. This includes drafts, books, sketches or training manuals. Necessary record keeping material will be made available at the work site location by the team judges.

NOTE the following equipment will be provided by the KDMM:

- Manikins
- KED or equivalents
- Long spine boards with straps
- Oxygen cylinders
- Oxygen masks with tubing
- Oxygen regulator and wrench
- Traction splints
- Rigid splints

Problems will be designed from the Brady First Response text by Grant Goold and Scott Vahradian. The contest rules and skill sheets may also be subject to change by the Contest Committee.

Upon arrival at the contest field, the MET team will introduce themselves to the judges

^{*(1)} Aspirator, Hand Bulb with catheter

^{*(4)} Heat Packs

^{*(1)} Poison Antidote Kit

^{*(1)} Roll aluminum Foil, 18"

^{*(6)} Roller Bandage, Gauze, 4" x 5 yds.

^{*(2)} Splints, Wire Ladder or Equivalent

^{*} These items can be obtained from the MET/EMT Station.

and identify the primary and secondary MET. The team will place their patient in the designated area. The chairman of the judges will furnish a copy of the hearsay statement to the primary MET after the clock has been started. The MET team will not be allowed to lay out their equipment or don any infectious control equipment or apparel until after the clock has started.

Problems may be designed with life-threatening injuries or medical problems that require immediate attention of the MET member(s), prior to systematic patient assessment/examination by the MET team.

Conditions, signs and symptoms associated with patient injuries or illnesses will be provided to the Primary MET by the contest judge.

Ventilation, cardiac arrest and oxygen therapy problems will be performed on manikins provided by the KDMM. Live patients will not be used for these skills.

All manikins shall be thoroughly cleaned after each use by the contest grounds crew. Alcohol preps and paper towels will also be furnished by the contest grounds crew.

Skill Evaluation Station

In addition to the skills performed during working of the problem, one or both METS may be required to perform one or more of the following skills at the Skill Evaluation Station:

Bag Valve Mask Ventilation - Skill Sheet Number 2
Oxygen Administration - Skill Sheet Number 4
Suctioning - Skill Sheet Number 9
Nasopharyngeal Airway - Skill Sheet Number 10
Mouth to Mask Ventilation - Skill Sheet Number 11
Bleeding Control, Bandaging, and Splinting - Skill Sheets No. 12 thru 13J
Two Man Direct Ground Lift - Skill Sheet Number 14
Adult 1-Rescuer CPR - Skill Sheet Number 15
Complete Airway Obstruction (Conscious Patient) - Skill Sheet Number 16
Airway Obstruction (Victim Loses Consciousness) - Skill Sheet Number 17
Obstructed Airway (Unconscious Victim-Unwitnessed) - Skill Sheet Number 18
Two Rescuer CPR - Skill Sheet Number 19
Establishing Airway , Suspected Cervical Injury - Skill Sheet Number 20

Note: BSI Precautions must be taken while performing the above skill(s) and total time will be recorded for completing skill(s).

Contest Judging

A judge appointed by the <u>Chief Judge</u> will be placed in the team lock-up area and may examine and check each team's equipment. Any items not found in accordance with requirements will result in team discounts.

At least two (2) judges will be assigned to each team while working the problem and an additional person may serve as timekeeper. However, one of the two judges can assume the timekeeper's duties if necessary. Both judges must be Mine Emergency Technicians (MET) as certified by the Kentucky Department of Mines and Minerals. One of the judges will serve as Chairman and will gather all data and material from the other judge upon completion of the problem.

Judges shall give full attention to teams during working of the problem but any unnecessary communication is prohibited. Judges shall not be affiliated with any of the participating teams or organizations.

The performance of MET teams while working problems will be judged by the use of MET contest skill sheets developed by the MET Contest Committee. These skill sheets will be provided to all interested persons prior to the first MET Contest. Any discounts will be noted on one or more of the following score cards:

Scorecard A------Recommended Functions and Critical Skills Scorecard B-------Skill Evaluation Station Scorecard C-------Written Examination Scorecard D------Time for Problem Completion

At least two judges must concur on the assessment of team discounts. <u>Judges will not discuss discounts with participating teams prior to the appeals process.</u>

NOTE: All contest material and skill sheets shall be signed by the judges and delivered to the score card examiner for reviewing.

Written Examination

The written examination shall consist of 20 or less multiple choice questions and/or "fill in the blank" taken from the Brady First Response text, by Grant Goold and Scott Vahradian. The questions may be taken from one or more of the textbook chapters. However, no questions will be taken from Chapter 24 or 30.

One or more judges shall administer the written exam.

Twenty minutes shall be allowed for taking the written examination. Judges are not allowed to explain the meaning of a question or any wording.

The Primary and Secondary MET from each participating team must take the written examination. Patients are not allowed to take written examinations.

While taking the written examination, team members will be separated in an effort to prevent discussion of questions and answers.

WRITTEN EXAMINATION

Rule 1 Written Exam: For each incorrect answer ------One (1) Discount.

GENERAL RULES DISCOUNTS

Rule 1 Failure to identify team members roles; Primary MET, Secondary MET and Patient (if provided)------Five (5) Discounts.

Prior to starting the contest clock, team member will declare to the contest judge, which member will serve as Primary and Secondary MET.

The Secondary MET may be directed by the Primary MET to assist in performing some of the recommended functions and critical skills. Example: While performing patient assessment on the leg, the Primary MET may be informed by the judge that profuse bleeding is occurring in the femur area. The Secondary MET may be instructed by the Primary MET to continue with patient leg assessment, but to not go beyond that recommended function, while the Primary MET takes action to stop the bleeding.

Rule 2 Failure to declare role change of team members to the contest judge, while performing problem (Example: Primary changes to Secondary MET)---Five (5) Discounts.

Team members may change from Primary to Secondary MET while working the problem, but the judge must be informed prior to the change taking place.

RULE 2 CLARIFICATION

When the Primary MET calls for a change from Primary to Secondary, the Secondary MET becomes the Primary MET and is responsible for performing recommended functions and critical skills. He then becomes responsible for directing the Secondary MET and must make the request for role reversal.

Rule 3 Failure to make an emergency move when indicated by a card from the judge ------Thirty (30) Discounts.

When working the problem, teams may be given cards by the judges stating life-threatening conditions or an unsafe patient location which will require an emergency move by the MET team.

Emergency moves used at ground level are the Shirt Drag, Blanket Drag and the Shoulder Drag.

condition-----Five (5) Discounts.

- **Rule 6** Any action by a team member that can be life-threatening to the patient-----Thirty (30) Discounts.
- **Rule 7** Failure to comply with required inventory for first response kit (one discount for each item missing)------One (1) discount.

Rule 8A Failure to use or wear infection control apparel while examining a patient-----Five (5) Discounts.

Rule 8B Failure to replace damaged infection control apparel (example: torn glove) while working the problem------Five (5) Discounts.

Rule 9 Failure of team to remove used and unused material from the contest work area within five (5) minutes of problem completion (stopping the clock)----Five (5) Discounts.

TIMING

Rule 1 Failure of MET team to complete problem in allotted time will result in discounts for recommended functions and critical skills not completed.

Rule 2 Performing work prior to starting the contest clock or after the clock has been stopped------Five (5) Discounts.

A maximum of 30 minutes will be allowed to complete the problem by the team, unless otherwise specified by the Chief Judge. Upon completion of the problem, the judges and the MET team participants will note the working time of the problem.

RECOMMENDED FUNCTIONS AND CRITICAL SKILLS

MET team members shall follow infection control procedures and wear infection control apparel while performing the problem.

Teams will be judged on performance of recommended functions and critical skills and the following discounts shall apply:

- Rule 1 Failure to perform a recommended function------Minimum: 5 Discounts

 Maximum: Equals total number of critical skills in function
- **Rule 2** Failure to perform a critical skill-----One (1) Discount.

Skill Evaluation Station

- **Rule 1** Failure to perform a recommended function-------Minimum: 5 Discounts Maximum: Equals total number of critical skills in function
- Rule 2 Failure to perform a critical skill------One (1) Discount.

See page 4 of this document for a list of skills that may have to be performed by one or both METS at the Skill Evaluation Station.

MINE EMERGENCY TECHNICIAN CONTEST SKILL SHEETS

PATIENT ASSESSMENT - UNCONSCIOUS PATIENT	PAGE 10
BAG-VALVE-MASK VENTILATION	
CERVICAL SPINE SPLINT (KED)	PAGE 12
OXYGEN ADMINISTRATIONCLOSED FEMUR FRACTURE - TRACTION SPLINT	PAGE 13
CLOSED FEMUR FRACTURE - TRACTION SPLINT	PAGE 14
THREE PERSON LOG ROLL_	PAGE 15
THREE PERSON LOG ROLL_ TWO RESCUER EXTREMITY GROUND LIFT	PAGE 16
SHIRT DRAG SUCTIONING USING THE NASOPHARYNGEAL AIRWAY	PAGE 17
SUCTIONING	PAGE 18
USING THE NASOPHARYNGEAL AIRWAY	PAGE 19
MOUTH-TO-MASK VENTILATION	PAGE 20
DRESSING AND BANDAGING	PAGE 21
BLEEDING CONTROL_	PAGE 22
SPLINTINGIMMOBILIZATION OF DISLOCATED HIP	PAGE 23
IMMOBILIZATION OF DISLOCATED HIP	PAGE 24
IMMOBILIZATION OF FRACTURED TIBIA OR FIBULA	PAGE 25
IMMOBILIZATION OF FRACTURED OR DISLOCATED ANKLE	
IMMOBILIZATION OF FRACTURED CLAVICLE	PAGE 27
IMMOBILIZATION OF DISLOCATED SHOULDER (ANTERIOR)	
IMMOBILIZATION OF FRACTURE TO HUMERUS	PAGE 29
IMMOBILIZATION OF DISLOCATED ELBOW	
USING A PADDED WIRE LADDER SPLINT	PAGE 30
IMMOBILIZATION OF FRACTURE TO FOREARM	
ONE INCH ABOVE THE WRIST	PAGE 31
AIR SPLINT RADIOUS AND ULNA	PAGE 32
TWO MAN DIRECT GROUND LIFT	PAGE 33
ADULT 1-RESCUER CPR	PAGE 34
COMPLETE AIRWAY OBSTRUCTION (CONSCIOUS PATIENT)	PAGE 35
AIRWAY OBSTRUCTION (VICTIM LOSES CONSCIOUSNESS)	PAGE 36
OBSTRUCTED AIRWAY UNCONSCIOUS VICTIM-UNWITNESSED_	PAGE 37
	PAGE 38
ESTABLISHING AIRWAY, SUSPECTED CERVICAL INJURY	PAGE 39

Note: Body substance isolation (BSI) precautions are not noted on some skill sheets, however BSI precautions must be taken while performing all skills.

PATIENT ASSESSMENT (MET) SKILL SHEET NUMBER 1

R. SCERS SIZES B. Mechanism if injury/nature of illness C. Number of patients D. Access and extrication needs E. Considers spinal immobilization A. Forms general expression B. Determine chief complain/apparent life threats C. Level of consciousness (AVPU) D. Assess airway E. Assess breathing/considers 02 therapy F. Assess sire color / temperature I. Identify patient priority and transport decision III. PHYSICAL EXAM AND TREATMENT VITALS: OBTAINS S.A.M.P.L.E. HISTORY: B. Pulse (rate and character) Note: Count pulse and respirations 30 seconds x 2 within 4 counts Signs and symptoms (PQRST) D. Allergies E. Medications F. Past pertinent history G. Last oral intake H. Events I. Head J. Neck (cervical spine/front and surface of neck) K. Medical alerts L. Scalp M. Eyes N. Ears O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert V. VERBALIZE PATIENT HAND OFF USING E.S.A.M.P.L. ACRONYM	I. SCENE SIZE-UP	A. Scene Safety	
II. INITIAL ASSESSMENT AND TREATMENT III. INITIAL ASSESSMENT AND TREATMENT III. INITIAL ASSESSMENT AND TREATMENT III. PHYSICAL EXAM AND TREATMENT VITALS: OBTAINS S.A.M.P.L.E. HISTORY: HEAD TO TOE EXAM USING D.O.T.S.: HEAD TO TOE EXAM USING D.O.T.S.: OBTAINS S.A.M.P.L.E. HISTORY: III. PHYSICAL EXAM AND TREATMENT VITALS: OBTAINS S.A.M.P.L.E. HISTORY: III. PHYSICAL EXAM AND TREATMENT VITALS: OBTAINS S.A.M.P.L.E. HISTORY: III. PHYSICAL EXAM AND TREATMENT VITALS: OBTAINS S.A.M.P.L.E. HISTORY: OBTAINS S.A.M.P.L.E. HISTORY: III. PHYSICAL EXAM USING D.O.T.S.: OBTAINS S.A.M.P.L.E. HISTORY: III. PHYSICAL EXAM AND TREATMENT VITALS: OBTAINS S.A.M.P.L.E. HISTORY: III. PHYSICAL EXAM AND TREATMENT PHICH PROPRIET TO	I. SCENE SIZE-UP		
II. INITIAL ASSESSMENT AND TREATMENT III. PHYSICAL EXAM AND TREATMENT VITALS: OBTAINS S.A.M.P.L.E. HISTORY: OBTAINS S.A.M.P.L.E. HISTORY: III. PHYSICAL EXAM USING D.O.T.S.: OBTAINS S.A.M.P.L.E. HISTORY: III. PHYSICAL EXAM USING D.O.T.S.: OBTAINS S.A.M.P.L.E. HISTORY: III. PHYSICAL EXAM OBTAINS S.A.M.P.L.E. HISTORY: OBTAINS S.A.M.P.L.E. HISTORY: OBTAINS S.A.M.P.L.E. HISTORY: III. PHYSICAL EXAM USING D.O.T.S.: OBTAINS S.A.M.P.L.E. HISTORY: OBTAINS S.A.M.P.L.E. HASTORY: OBTAINS M. ASSESS BRICH HISTORY: OBTAINS C. Level for complain and perbase solutions for the primary of the properties of the property of the proper			
II. INITIAL ASSESSMENT AND TREATMENT B. Determine chief complain/apparent life threats C. Level of consciousness (AVPU) D. Assess airway E. Assess breathing/considers 02 therapy F. Assess circulation / checks pulse G. Assess and control severe bleeding H. Assess sirculation / checks pulse G. Assess and control severe bleeding H. Assess sirculation / checks pulse G. Assess and control severe bleeding H. Assess skin color / temperature I. Identify patient priority and transport decision III. PHYSICAL EXAM AND TREATMENT VITALS: OBTAINS S.A.M.P.L.E. HISTORY: D. A. BP (+ or - 10 mmhg) B. Pulse (rate and character) C. Respirations (rate and character) C. Respirations (rate and character) C. Respirations (rate and character) C. Allergies E. Medications F. Past pertinent history G. Last oral intake H. Events I. Head J. Neck (cervical spine/front and surface of neck) K. Medical alerts L. Scalp M. Eyes N. Ears O. Nose P. Mouth Q. Face R. C.HEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe			
II. INITIAL ASSESSMENT AND TREATMENT A. Forms general expression B. Determine chief complain/apparent life threats C. Level of consciousness (AVPU) D. Assess airway E. Assess breathing/considers 02 therapy F. Assess sirculation / checks pulse G. Assess and control severe bleeding H. Assess skin color / temperature I. Identify patient priority and transport decision III. PHYSICAL EXAM AND TREATMENT VITALS: OBTAINS S.A.M.P.L.E. HISTORY: HEAD TO TOE EXAM USING D.O.T.S.: Note: Count pulse and respirations 30 seconds x 2 within 4 counts Signs and symptoms (PQRST) D. Allergies D. Allergies D. Medications F. Past pertinent history G. Last oral intake H. Events H.			
TREATMENT B. Determine chief complain/apparent life threats C. Level of consciousness (AVPU) D. Assess airway E. Assess breathing/considers 02 therapy F. Assess circulation / checks pulse G. Assess and control severe bleeding H. Assess skin color / temperature I. Identify patient priority and transport decision A. BP (+ or - 10 mmhg) B. Pulse (rate and character) C. Respirations (rate and character) Note: Count pulse and respirations 30 seconds x 2 within 4 counts Signs and symptoms (PQRST) D. Allergies E. Medications F. Past pertinent history G. Last oral intake H. Events I. Head J. Neck (cervical spine/front and surface of neck) K. Medical alerts L. Scalp M. Eyes N. Ears O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT	II INITIAI ASSESSMENT AND	1	-
C. Level of consciousness (AVPU) D. Assess airway E. Assess breathing/considers 02 therapy F. Assess circulation / checks pulse G. Assess and control severe bleeding H. Assess skin color / temperature I. Identify patient priority and transport decision III. PHYSICAL EXAM AND TREATMENT VITALS: OBTAINS S.A.M.P.L.E. HISTORY: B. Pulse (rate and character) C. Respirations (rate and character) Note: Count pulse and respirations 30 seconds x 2 within 4 counts Signs and symptoms (PQRST) D. Allergies E. Medications F. Past pertinent history G. Last oral intake H. Events I. Head J. Neck (cervical spine/front and surface of neck) K. Medical alerts L. Scalp M. Eyes N. Ears O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT			
D. Assess airway E. Assess breathing/considers 02 therapy F. Assess breathing/considers 02 therapy F. Assess and control severe bleeding H. Assess skin color / temperature I. Identify patient priority and transport decision A. BP (+ or - 10 mmhg) B. Pulse (rate and character) C. Respirations (rate and character) Note: Count pulse and respirations 30 seconds x 2 within 4 counts Signs and symptoms (PQRST) D. Allergies E. Medications F. Past pertinent history G. Last oral intake H. Events I. Head J. Neck (cervical spine/front and surface of neck) K. Medical alerts L. Scalp M. Eyes N. Ears O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe	TREATMENT		_
E. Assess breathing/considers 02 therapy F. Assess circulation / checks pulse G. Assess and control severe bleeding H. Assess skin color / temperature I. Identify patient priority and transport decision III. PHYSICAL EXAM AND TREATMENT VITALS: OBTAINS S.A.M.P.L.E. HISTORY: OBTAINS S.A.M.P.L.E. HISTORY: B. Pulse (rate and character) C. Respirations (rate and character) Note: Count pulse and respirations 30 seconds x 2 within 4 counts Signs and symptoms (PQRST) D. Allergies E. Medications F. Past pertinent history G. Last oral intake H. Events I. Head J. Neck (cervical spine/front and surface of neck) K. Medical alerts L. Scalp M. Eyes N. Ears O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe		, ,	_
F. Assess circulation / checks pulse G. Assess and control severe bleeding H. Assess skin color / temperature I. Identify patient priority and transport decision III. PHYSICAL EXAM AND TREATMENT VITALS: OBTAINS S.A.M.P.L.E. HISTORY: B. Pulse (rate and character) C. Respirations (rate and character) C. Respirations (rate and character) Note: Count pulse and respirations 30 seconds x 2 within 4 counts Signs and symptoms (PQRST) D. Allergies E. Medications F. Past pertinent history G. Last oral intake H. Events I. Head J. Neck (cervical spine/front and surface of neck) K. Medical alerts L. Scalp M. Eyes N. Ears O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe		·	_
G. Assess and control severe bleeding H. Assess skin color / temperature I. Identify patient priority and transport decision A. BP (+ or - 10 mmhg) TREATMENT VITALS: OBTAINS S.A.M.P.L.E. HISTORY: B. Pulse (rate and character) C. Respirations (rate and character) Note: Count pulse and respirations 30 seconds x 2 within 4 counts Signs and symptoms (PQRST) D. Allergies E. Medications F. Past pertinent history G. Last oral intake H. Events I. Head J. Neck (cervical spine/front and surface of neck) K. Medical alerts L. Scalp M. Eyes N. Ears O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT			
H. Assess skin color / temperature I. Identify patient priority and transport decision A. BP (+ or - 10 mmhg) TREATMENT VITALS: OBTAINS S.A.M.P.L.E. HISTORY: HEAD TO TOE EXAM USING D.O.T.S.: Note: Count pulse and character) Note: Count p			
III. PHYSICAL EXAM AND TREATMENT VITALS: OBTAINS S.A.M.P.L.E. HISTORY: OBTAIN S D.O.T.S.: HEAD TO TOE EXAM USING D.O.T.S.: HEAD TO TOE EXAM		-	
III. PHYSICAL EXAM AND TREATMENT VITALS: OBTAINS S.A.M.P.L.E. HISTORY: Note: Count pulse and respirations 30 seconds x 2 within 4 counts Signs and symptoms (PQRST) D. Allergies E. Medications F. Past pertinent history G. Last oral intake H. Events I. Head J. Neck (cervical spine/front and surface of neck) K. Medical alerts L. Scalp M. Eyes N. Ears O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT			
B. Pulse (rate and character) C. Respirations (rate and character) Note: Count pulse and respirations 30 seconds x 2 within 4 counts Signs and symptoms (PQRST) D. Allergies E. Medications F. Past pertinent history G. Last oral intake H. Events I. Head J. Neck (cervical spine/front and surface of neck) K. Medical alerts L. Scalp M. Eyes N. Ears O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT		I. Identify patient priority and transport decision	
OBTAINS S.A.M.P.L.E. HISTORY: Note: Count pulse and respirations 30 seconds x 2 within 4 counts Signs and symptoms (PQRST) D. Allergies E. Medications F. Past pertinent history G. Last oral intake H. Events I. Head J. Neck (cervical spine/front and surface of neck) K. Medical alerts L. Scalp M. Eyes N. Ears O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT	III. PHYSICAL EXAM AND	A. BP (+ or – 10 mmhg)	
OBTAINS S.A.M.P.L.E. HISTORY: Note: Count pulse and respirations 30 seconds x 2 within 4 counts Signs and symptoms (PQRST) D. Allergies E. Medications F. Past pertinent history G. Last oral intake H. Events I. Head J. Neck (cervical spine/front and surface of neck) K. Medical alerts L. Scalp M. Eyes N. Ears O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT	TREATMENT VITALS:	B. Pulse (rate and character)	
HISTORY: Within 4 counts Signs and symptoms (PQRST) D. Allergies E. Medications F. Past pertinent history G. Last oral intake H. Events I. Head J. Neck (cervical spine/front and surface of neck) K. Medical alerts L. Scalp M. Eyes N. Ears O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT			
HEAD TO TOE EXAM USING D.O.T.S.: Signs and symptoms (PQRST) D. Allergies E. Medications F. Past pertinent history G. Last oral intake H. Events I. Head J. Neck (cervical spine/front and surface of neck) K. Medical alerts L. Scalp M. Eyes N. Ears O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT IV. VERBALIZE NEED FOR ONGOI		Note: Count pulse and respirations 30 seconds x 2	
D. Allergies E. Medications F. Past pertinent history G. Last oral intake H. Events I. Head J. Neck (cervical spine/front and surface of neck) K. Medical alerts L. Scalp M. Eyes N. Ears O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT Interval in the story I. Medications E. Medications	HISTORY:	within 4 counts	
HEAD TO TOE EXAM USING D.O.T.S.: E. Medications F. Past pertinent history G. Last oral intake H. Events I. Head J. Neck (cervical spine/front and surface of neck) K. Medical alerts L. Scalp M. Eyes N. Ears O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT		Signs and symptoms (PQRST)	
HEAD TO TOE EXAM USING D.O.T.S.: F. Past pertinent history G. Last oral intake H. Events I. Head J. Neck (cervical spine/front and surface of neck) K. Medical alerts L. Scalp M. Eyes N. Ears O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT			
D.O.T.S.: G. Last oral intake H. Events I. Head J. Neck (cervical spine/front and surface of neck) K. Medical alerts L. Scalp M. Eyes N. Ears O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT		E. Medications	
D.O.T.S.: G. Last oral intake H. Events I. Head J. Neck (cervical spine/front and surface of neck) K. Medical alerts L. Scalp M. Eyes N. Ears O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT	HEAD TO TOE EXAM USING	F. Past pertinent history	
H. Events I. Head J. Neck (cervical spine/front and surface of neck) K. Medical alerts L. Scalp M. Eyes N. Ears O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT		G. Last oral intake	
J. Neck (cervical spine/front and surface of neck) K. Medical alerts L. Scalp M. Eyes N. Ears O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT	D.O.1.5	H. Events	
K. Medical alerts L. Scalp M. Eyes N. Ears O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT		I. Head	
L. Scalp M. Eyes N. Ears O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT		J. Neck (cervical spine/front and surface of neck)	
M. Eyes N. Ears O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT		K. Medical alerts	
N. Ears O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT		L. Scalp	
O. Nose P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT		M. Eyes	
P. Mouth Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT		N. Ears	
Q. Face R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT		O. Nose	
R. CHEST: Collar bone S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT		P. Mouth	
S. Ribs and chest wall T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT		Q. Face	
T. Signs of respiratory distress Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT			
Abdomen and Pelvis: U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT		S. Ribs and chest wall	
U. Abdomen surface V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT			
V. Pelvis W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT			
W. Legs, Arms: Pulses, motor sensory, medical alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT			
alert X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT			
X. Back surfaces: Examines back from head to toe IV. VERBALIZE NEED FOR ONGOING ASSEMENT			
IV. VERBALIZE NEED FOR ONGOING ASSEMENT			
IV. VERBALIZE NEED FOR ONGOING ASSEMENT			
	IV. VERBALIZE NEED FOR ONGOIN		

BAG VALVE-MASK VENTILATION (MET) SKILL SHEET NUMBER 2

RECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
1. BSI	Take BSI precautions		
2. SCENE SIZE-UP	Performs scene size-up		
3. DETERMINE	Determine unresponsiveness		
UNRESPONSIVENESS			
4. AIRWAY-BREATHING	Open patient's airway and assess breathing		
5. VENTILATE PATIENT	Ventilate patient with two breaths		
6. CHECK CIRCULATION	Check circulation/carotid pulse		
7. VENTILATION PATIENT	Ventilate patient every 5 seconds (while		
	measuring and inserting OPA and		
	assembling BVM)		
8. SELECT OPA	Determine the proper size OPA (between		
	ventilations)		
9. INSERT OPA	Insert the OPA upside down		
10. ROTATE OPA	Rotate the OPA 180 degrees clockwise		
	until the flange rests on patient's teeth		
	(between ventilations)		
11. REASSESS AIRWAY AND	Reassess the patient's airway and		
BREATHING	breathing		
12. SELECT BVM	Select the correct size BVM for the patient		
	(between ventilations)		
13. POSITION MASK	Position the mask, creating a proper mask-		
	to-face seal, make sure the bag is		
	connected to the mask.		
14. VENTILATE PATIENT	Squeeze the bag until the patient's chest rises.		
	(If you are alone, compress the bag against		
	your thigh. If you are with a partner, hold the		
	seal while your partner compresses the bag		
	with two hands.)		
	NOTE: Ventilate the patient, making sure		
	the chest rises and that you hear air		
	escaping during exhalationIf your patient		
	gags, remove the OPA		

BSI - Body Substance Isolation

BVM - Bag Valve Mask

OPA - Oropharyngeal Airway

RECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
1. POSITIONING	A. M.E.T. 2 – Take stability position and on		
	command from M.E.T. 1, move patient into		
	proper position and maintain support until		
	chest straps are fastened and snug		
2. APPLICATION OF	A. Apply extrication collar		
EXTRICATION DEVICE BY	B. Slide device behind patient and into proper		
MET 1	place (centered) and snug under arm pits		
	C. Fasten bottom two straps (red and yellow)		
	D. Place leg straps, pass straps around thigh and		
	fasten on same side with thighs padded		
	E. Pad behind head and fasten forehead strap		
	F. Fasten top chest strap (green)		
	G. Ensure all three front straps are snug		
3. LIFTING	A. Both M.E.T.'s lift under armpits and thighs		
	and transfer patient to stretcher or long board		

RECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
1. BSI	Take or verbalize BSI precautions		
2. AIRWAY MAINTENANCE	Maintain patient's airway and breathing		
3. CHECK OXYGEN SOURCE	Check for medical 0 ₂ while maintaining stability of		
	tank		
4. SEAL REMOVAL	Remove the protective seal from the tank		
5. CLEAN VALVE	Open and shut valve quickly to clean away debris		
6. REGULATOR ASSEMBLY	Assemble regulator-flowmeter to tank		
7. OPEN TANK VALVE	When regular is hand-tight, open tank valve by		
	making one-quarter turn		
8. ATTACHED MASK OR	Attach the proper delivery device (non-rebreather		
NASAL CANNULA	mask or nasal cannula) to the flowmeter		
9. ADJUST FLOWMETER	Open the flowmeter to the desired setting (non-		
	rebreather mask – 15 liters per minute and verify		
	bag is filled: nasal cannula – 6 liter per minute)		
10. DELIVERY DEVICE	Apply delivery device to the patient and administer		
APPLICATION (MASK OR	oxygen		
NASAL CANNULA)			
11. PREVENTION OF BLOOD	Prevent any further blood loss		
LOSS			
12. POSITION PATIENT	Properly position the patient (shock position unless		
12 COLUED DA EVENTE	contraindicated)		
13. COVER PATIENT	Cover the patient with a blanket		
14. TRANSPORT	Arrangement for immediate transport		

BSI – BODY SUBSTANCE ISOLATION

CLOSED PAINFUL SWOLLEN DEFORMED INJURY TO MID THIGH-TRACTION SPLINT

(MET) Skill Sheet Number 5

R	ECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
1.	CIRCULATION CHECK	Perform circulation check		

2. SENSATION CHECK	Perform sensation check and check contraindications Caution: Do not use the traction splint with the following injuries: The injury is an open type injury. The knee of the injured leg is also injured. The injury is close to	
	the knee. The lower leg or ankle is injured. The hip is injured. The pelvis is injured. The lower leg is partially amputated.	
3. STABILIZE AND APPLY MANUAL TRACTION	Stabilize the injured leg and apply manual traction when required. Do not release it until the leg is completely immobilized. (The ankle hitch may be attached before manual traction is applied with some types of splinting).	
4. PREPARE AND ADJUST SPLINT	Prepare and adjust splint to proper length using the uninjured leg as a guide, then place it under the injured leg	
5. FASTEN ISCHIAL STRAP	Fasten ischial strap	
6. APPLY ANKLE HITCH	Apply ankle hitch if not already attached	
7. APPLY MECHANICAL TRACTION	Apply mechanical traction to the splint	
8. RELEASE MANUAL TRACTION	Release manual traction when mechanical traction is adequate	
9. FASTEN STRAPS OR CRAVATS	Fasten straps or cravats to adequately secure leg to splint	
10. RE-ASSESS SPLINT	Re-assess splint for adequate traction	
11. PERFORM CIRCULATION CHECK	Perform circulation check	
12. PERFORM SENSATION CHECK	Perform sensation check	

THREE-PERSON LOG ROLL (MET) Skill Sheet Number 6

RECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
1. STABILIZE HEAD	A. M.E.T. #1 will stabilize the head and neck and		
	will direct the others to roll the patient as a unit,		
	on command at appropriate time.		
	B. M.E.T. #2 will set collar in place		

	C. M.E.T. #2 will secure collar	
2. PREPARING THE	A. M.E.T. #3 will place a long backboard parallel to	
PATIENT	the patient	
	B. M.E.T. #2 will kneel at the level of the victim's	
	shoulders on the side opposite the board leaving	
	room to roll the patient toward knees	
	C. If the patient's arm nearest the rescuer is not	
	injured, the second M.E.T. will raise the arm	
	above the patient's head.	
3. POSITIONING OF	A. M.E.T. #2 grasps the patient at the shoulder and	
RESCUERS	hip areas	
	B. M.E.T. #3 will grasp the patient above the knees	
	and near the ankles (Met #1 maintains head	
	stabilization)	
4. ROLLING THE PATIENT	A. On command from the M.E.T. stabilizing the	
	head, the patient is rolled toward the rescuer by	
	pulling steadily and evenly at the shoulder and	
	pelvis area.	
	B. The head and neck should remain on the same	
	plane as the torso	
	C. M.E.T. #2 must maintain stability by holding	
	patient with one hand and positioning board with	
	other.	
	D. Roll the patient as a unit onto the board (board	
	may be slanted or flat)	
	E. Place the patient's arm alongside him.	
5. SECURING PATIENT TO	A. M.E.T. #2 will secure the patient's body (torso,	
BOARD	legs) to the already prepared board while head	
	support is maintained	
	B. M.E.T. #2 will secure the patient's head to the	
	Board	
	C. M.E.T. #2 will tie the wrists together loosely	
	unless injury prevents	

TWO RESCUER EXTREMITY GROUND LIFT (MET) Skill Sheet Number 7

RECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
1. POSITIONING	A. M.E.T. 1 – kneel at the head of the patient and		
	place one hand under each of the shoulders		
	B. M.E.T. 2 – kneel by the patient's knees and		
	grasp the patient's wrists		

2. RAISING PATIENT	A. M.E.T. 1 – push patient's shoulders up and support patient's back and head with body.	
	B. M.E.T. 2 – gently pull on patient's arms, raising to a sitting position	
3. POSITIONING AND LIFTING	A. M.E.T. 1 – support patient in sitting position and slip hands under patient's arms and grasp patient's wrists	
	B. M.E.T. 2 – slip hands under the patient's knees	
	C. On M.E.T.1 command, M.E.T.s stand	
	simultaneously, lifting patient with proper body mechanics	

SHIRT DRAG (MET) Skill Sheet Number 8

RECOMMENDED FUNCTION	Y	N
1. M.E.T. states need for emergency move		
2. M.E.T. takes position at victim's head		
3. M.E.T. grasps shirt at the shoulder area		
4. M.E.T. drags patient in straight line		

_

SUCTIONING (MET) Skill Sheet Number 9

RECOMMENDED FUNCTION CRITICAL SKILL Take or verbalize BSI Precautions 1. BSI 2. TURN DEVICE ON Prepare and turn on the suction device (or equivalent) Assure presence of mechanical suction 3. CHECK DEVICE 4. INSERT CATHETER Without suction, insert the suction catheter to the base of patient's tongue 5. BEGIN SUCTIONING Begin suctioning while moving the tonsil tip from side to side as you remove it from the mouth 6. SUCTION AND VENTILATE Suction for up to 15 seconds in an adult, then ventilate for two minutes if patient is not breathing; suction again if necessary NOTE: If the suction catheter becomes clogged, rinse it out and continue

RECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
1. BSI	Take or verbalize BSI Precautions		
2. NPA SIZE	Determine the proper size NPA (measure from tip of		
	nose to tip of ear		
3. LUBRICATE NPA	Lubricate the NPA with a water-soluble lubricant		
4. POSITION NPA	Position the NPA with the beveled end toward the		
	septum		
5. ADVANCE NPA	Advance the NPA gently with a rotating motion		
6. REASSESS PATIENT	Reassess the patient to ensure an open airway and		
	breathing NOTE: if you meet resistance during		
	insertion, try the same procedure on the other nostril		

BSI – BODY SUBSTANCE ISOLATION

NPA – NASOPHARYNGEAL AIRWAY

RECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
1. PERFORM SCENE SIZE-UP	A. Takes BSI precautions		
	B. Checks scene safety		
	C. Mechanism of injury/nature of illness		
	D. Number of patients		
	E. Access and extrication needs		
2. ASSESS LEVEL OF	A. Shouts "Are you OK?"		
CONSCIOUSNESS	B. Taps or gently shakes shoulders (does not compromise c-spine)		
	C. Determine unresponsiveness. Activates EMS System		
3. ESTABLISH AIRWAY	Correctly executes head-tilt / chin-lift or jaw-		
4. CHECK PATIENT'S	thrust maneuver depending on c-spine injury		
BREATHING	A. Look, listen, and feel for air exchange		
DREATHING	B. Determines absence of breathing 3-5 seconds		
5. VENTILATES PATIENT	C. Maintains airway		
5. VENTILATES PATIENT	A. Place and seal mask on patient		
	B. Ventilates patient 2 times C. Maintains airway		
	D. Considers 0 ₂ therapy		
6. CHECKS FOR CAROTID	A. Correctly locate area of carotid pulse		
PULSE	B. Slides 2 or 3 fingers into groove on same side of		
TOLSE	neck as rescuer		
	C. Maintain open airway and indicate presence of pulse		
7. CONTINUE	A. Determines need to continue ventilations only		
VENTILATIONS	B. Maintain open airway		
	C. Maintain adequate seal with pocket mask		
	D. Continue ventilations for at least one minute		
	E. Deliver one breath every 5 or 6 seconds (10-12		
8. EVALUATE PULSE	breaths per minute)		
o. EVALUATE PULSE	A. Periodically (at least twice a minute) evaluates continued presence of pulse		

BSI – BODY SUBSTANCE ISOLATION

DRESSING, BANDAGING, AND BLEEDING CONTROL (MET) Skill Sheet Number 12-A

RECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
1. Proper application of dressing and bandage f	For top of head		
2. Proper application of dressing and bandage f	For burn injury on hand		
3. Proper application of dressing and bandage f	For cheek or jaw		
4. Proper application of dressing and bandage f	for protruding eyeball		
5. Proper application of dressing and bandage f	For hip or shoulder		
6. Proper application of dressing and bandage f	For sucking chest wound		
7. Proper application of dressing and bandage f	For impaled object in arm or leg		
8. Proper application of dressing and bandage f	For knee or elbow		
9. Proper application of dressing and bandage f	for the neck		
Each Performance Should	A. Selection of appropriate dressing		
Include:	and bandage		
metude.	B. Sufficient wound coverage		
	C. Bandage secure		
	D. No loose ends		
	E. Checking circulation distal to injury		
	when appropriate		

BLEEDING CONTROL (MET) Skill Sheet Number 12-B

RECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
1. Take or verbalize BSI Precautions			
2. Apply direct pressure to the bleeding wour	nd when appropriate		
3. Elevate the extremity, but only if there is n	o major injury to the underlying muscle or		
bone			
4. Reassess the wound; if bleeding does not s	top, apply additional pressure as needed		
5. If bleeding continues, apply pressure to the on the injured extremity	e arterial pressure point above the wound		
6. When bleeding is under control, dress and	bandage appropriately		
7. EMERGENCY CARE FOR AN OPEN	A. Expose Wound		
WOUND	B. Clear wound surface		
	C. Continue to control bleeding		
	D. Prevent further contamination		
	E. Keep patient lying still		
	F. Reassure the patient		
	G. Treat for shock		
8. APPLY DRESSING	A. Use sterile dressing		
	B. Cover entire wound		
	C. Control bleeding		
	D. Do not remove dressing		
9. APPLY BANDAGE	A. Bandage dressing in place after		
	bleeding is controlled		
	B. Do not bandage too tightly		
	C. Do not bandage too loosely		
	D. Do not leave loose ends		
	E. Do not cover the tips of fingers or		
	toes appropriate		
	F. Cover all edges of dressing		

BSI – BODY SUBSTANCE ISOLATION

RECOMMENDED FUNCTION

Y N

1. Take or verbalize BSI precautions	
2. Apply manual stabilization to the injured extremity	
3. Assess pulse, sensation, and movement below the injury site	
4. Measure the splint, and pad it appropriately	
5. For a long-bone injury, apply the splint so that the joints above and below the injury site are immobilized. For a joint injury, apply the splint so that the bones above and below the joint are	
immobilized.	
6. Secure the injured extremity to the splint	
7. Reassess pulse, sensation, and movement in the extremity. Document findings	
NOTE: If there is a deformity, and if the extremity below the injury is cyanotic or has	
no pulse, then align the extremity with gentle traction. However, if the injury site is at a	
joint, stop traction immediately if you feel any resistance.	

BSI – BODY SUBSTANCE ISOLATION

SPLINTING SKILLS

- 1. Immobilization of dislocated hip
- 2. Immobilization of fractured tibia or fibula
- 3. Immobilization of fractured or dislocated ankle
- 4. Immobilization of fractured clavicle
- 5. Immobilization of dislocated shoulder
- 6. Immobilization of fractured humerus
- 7. Immobilization of dislocated elbow
- 8. Immobilization of fractured forearm one inch above the wrist
- 9. Immobilization of fractured radius or ulna

SPLINTING OF LOWER EXTREMITIES (FRACTURES AND DISLOCATIONS) IMMOBILIZATION OF DISLOCATED HIP

RECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
1. IDENTIFICATION OF HIP	A. Anterior hip dislocation		
DISLOCATION	•		
	B. Lower limb is rotated outward		
	C. Hip is usually flexed		
	D. Posterior hip dislocation		
	E. Patient's leg rotated inward		
	F. The hip is flexed		
	G. The knee is bent		
	H. Foot may hang loose (foot drop)		
2. CARE FOR HIP	A. Check for distal pulse		
DISLOCATION			
	B. Check for sensation		
	C. Place folded blanket between patient's legs		
	D. Bind legs together with cravats		
3. SECURING PATIENT TO	A. Move patient to long spine board, use		
LONG SPINE BOARD	scoop style stretcher if available		
	B. Immobilize limb with pillows or rolled		
	blankets		
	C. Secure the patient to board with straps or		
	cravats		
	D. Reassess pulse and sensation		

SPLINTING OF LOWER EXTREMITY (FRACTURES AND DISLOCATIONS)

IMMOBILIZATION OF FRACTURED TIBIA OR FIBULA (USING AIR SPLINT)

CRITICAL SKILL	Y	N
A. Obvious signs of deformity (Often absent)		
B. Obvious signs of swelling and discoloration		
C. Conscious patient check for pain and		
tenderness		
A. Check for distal pulse		
B. Check for sensation		
C. Immobilize fracture with air-inflated splint		
A. Grasp leg with one hand above the injury		
site and other hand below injury site		
B. Maintain support		
C. Properly apply splint		
D. Splint should be relatively free of		
wrinkles		
E. Check for distal pulse prior to inflation		
F. Inflate splint to point that slight dent can be		
made		
G. Check distal pulse after inflation		
A. Check for leaks		
B. Periodically check for increase or decrease		
in pressure		
C. Monitor pressure in splint with finger tip		
D. Make certain desired pressure is		
maintained		
	 A. Obvious signs of deformity (Often absent) B. Obvious signs of swelling and discoloration C. Conscious patient check for pain and tenderness A. Check for distal pulse B. Check for sensation C. Immobilize fracture with air-inflated splint A. Grasp leg with one hand above the injury site and other hand below injury site B. Maintain support C. Properly apply splint D. Splint should be relatively free of wrinkles E. Check for distal pulse prior to inflation F. Inflate splint to point that slight dent can be made G. Check distal pulse after inflation A. Check for leaks B. Periodically check for increase or decrease in pressure C. Monitor pressure in splint with finger tip D. Make certain desired pressure is 	 A. Obvious signs of deformity (Often absent) B. Obvious signs of swelling and discoloration C. Conscious patient check for pain and tenderness A. Check for distal pulse B. Check for sensation C. Immobilize fracture with air-inflated splint A. Grasp leg with one hand above the injury site and other hand below injury site B. Maintain support C. Properly apply splint D. Splint should be relatively free of wrinkles E. Check for distal pulse prior to inflation F. Inflate splint to point that slight dent can be made G. Check distal pulse after inflation A. Check for leaks B. Periodically check for increase or decrease in pressure C. Monitor pressure in splint with finger tip D. Make certain desired pressure is

SPLINTING OF LOWER EXTREMITY (FRACTURES AND DISLOCATIONS)

IMMOBILIZATION OF FRACTURED OR DISLOCATED ANKLE

RECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
1. IDENTIFICATION OF	A. Obvious signs of deformity		
FRACTURED OR DISLOCATED ANKLE	B. Obvious signs of swelling and discoloration		
	C. Conscious patient check for pain and tenderness		
2. CARE FOR FRACTURED OR	A. Check for distal pulse		
DISLOCATED ANKLE	B. Check for sensation		
	C. Immobilize fracture or dislocation with pillow and cravats		
3. IMMOBILIZE FRACTURE OR DISLOCATION	A. Stabilize limb, lift limb, but do not apply traction		
	B. Place three cravats under ankle		
	C. Place pillow length wise under ankle, on top of cravats (pillow should extend 6 inches beyond foot)		
	D. Lower limb, adjust cravats for tying		
	E. Tie cravats distal to proximal		
	F. Tie fourth cravat at arch of foot		
	G. Elevate with blanket or pillow		
	H. Reassess distal pulse and sensation		

DISTAL: Means farther away from the torso

PROXIMAL: Means closer to the torso

SPLINTING OF UPPER EXTREMITIES (FRACTURES AND DISLOCATIONS)

IMMOBILIZATION OF FRACTURED CLAVICLE

RECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
1. IDENTIFICATION OF	A. Obvious signs of deformity		
FRACTURED CLAVICLE	B. Obvious signs of swelling and discoloration		
	C. Check for pain and tenderness		
2. CARE FOR FRACTURED CLAVICLE	A. Check for distal pulse		
	B. Check for sensation		
	C. Immobilize fracture with a sling and swathe		
3. IMMOBILIZING FRACTURE	A. Place sling over patients chest and under arm		
	B. MET 2 should hold or stabilize arm		
	C. Triangle should extend behind elbow on injured side		
	D. Pull sling around neck until hand is elevated and tie on uninjured side		
	E. Secure excess material at patients elbow		
	F. Patients fingertips should be exposed		
	G. Check for pulse and nerve function		
4. SECURING SLING WITH SWATHE	A. Use triangle cravat		
	B. Swathe is tied around chest and injured arm		
	C. Reassess distal pulse		
	D. Reassess sensation		
	E. Treat for shock		

MINE EMERGENCY TECHNICIAN (MET) Skill Sheet No. 13 - E

SPLINTING OF UPPER EXTREMITY

(FRACTURES AND DISLOCATIONS)

IMMOBILIZATION OF DISLOCATED SHOULDER (ANTERIOR)

RECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
1. IDENTIFICATION OF	A. Obvious signs of deformity		
DISLOCATED SHOULDER	B. Obvious signs of swelling and discoloration		
	C. Check for pain and tenderness		
2. CARE FOR DISLOCATED SHOULDER	A. Check for distal pulse		
	B. Check for sensation		
	C. Immobilize dislocation with appropriate_		
	padding, sling and swathe		
3. IMMOBILIZING DISLOCATED SHOULDER	A. Place appropriate padding between arm and chest		
	B. Place sling over appropriate padding and rest arm in position		
	C. MET 2 should hold or stabilize arm		
	D. Triangle should extend behind elbow on injured side		
	E. Pull sling around neck until hand is elevated and tie on uninjured side		
	F. Secure excess material at patients elbow		
	G. Patients fingertips should be exposed		
	H. Check for pulse and nerve function		
4. SECURING SLING WITH SWATHE	A. Use triangle cravat		
	B. Swathe is tied around chest and injured arm		
	C. Reassess distal pulse		
	D. Reassess sensation		
	E. Treat for shock		

MINE EMERGENCY TECHNICIAN (MET) Skill Sheet No. 13 - F

SPLINTING OF UPPER EXTREMITY (FRACTURES AND DISLOCATIONS)

IMMOBILIZATION OF FRACTURED HUMERUS

RECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
1. IDENTIFICATION OF	A. Obvious signs of deformity proximal, distal		
FRACTURED HUMERUS	or midshaft of humerus		
	B. Obvious signs of swelling and discoloration		
	C. Check for pain and tenderness		
2. CARE FOR FRACTURED HUMERUS	A. Check for distal pulse		
	B. Check for sensation		
	C. Immobilize fractured humerus with sling and swathe		
3. IMMOBILIZING A FRACTURED	A. Place sling over patients chest and place		
HUMERUS	arm over sling		
	B. MET 2 should hold or stabilize arm at a 90 degree angle		
	C. Triangle should extend behind elbow on injured side except for midshaft fracture		
	D. Pull sling around neck until hand is at a 90 degree angle and tie on uninjured side		
	E. Secure material at patients elbow if full		
	sling is used		
	F. Patients fingertips should be exposed		
	G. Check for pulse and nerve function		
4. SECURING SLING WITH SWATHE	A. Use triangle cravat		
	B. Swathe is tied around chest and injured arm		
	C. Reassess distal pulse		
	D. Reassess sensation		
	E. Treat for shock		

SPLINTING OF UPPER EXTREMITY FRACTURES AND DISLOCATIONS

IMMOBILIZATION OF A SUSPECTED DISLOCATION OF THE ELBOW, USING A PADDED WIRE LADDER SPLINT

RECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
1. IDENTIFICATION OF	A. Obvious signs of deformity		
DISLOCATED ELBOW	B. Obvious signs of swelling and discoloration		
	C. Check for pain and tenderness		
2. CARE FOR A DISLOCATED ELBOW	A. Check for distal pulse		
	B. Check for sensation		
	C. Immobilize dislocated elbow using a padded wire ladder splint		
3. IMMOBILIZING A DISLOCATED ELBOW	A. Prepare wire ladder splint with appropriate padding		
	B. Shape wire padded ladder splint with deformity of elbow		
	C. Stabilize limb, lift limb, but do not apply traction		
	D. Properly apply splint with appropriate wrap		
	E. Apply wrap distal to proximal		
	F. Check for distal pulse		
	G. Check for sensation		
4. IMMOBILIZING A DISLOCATED ELBOW FOR TRANSPORT	A. If elbow is dislocated in a bent position apply a wrist sling and swathe if possible		
	B. If elbow is dislocated in a straight position, place padding between splinted arm and body		
	C. Secure splinted limb to body with two cravats		
	D. Avoid placing cravats over injury site		
	E. Reassess pulse and sensation		

MINE EMERGENCY TECHNICIAN (MET) Skill Sheet No. 13 - H

SPLINTING OF UPPER EXTREMITY FRACTURES AND DISLOCATIONS

IMMOBILIZATION OF A SUSPECTED FRACTURE IN THE FOREARM 1 INCH ABOVE THE WRIST

USING A SHORT RIGID SPLINT, KERLIX OR CRAVATS AND SLING

RECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
1. IDENTIFICATION OF	A. Obvious signs of deformity		
FRACTURE			
WRIST AND FOREARM AREA	B. Obvious signs of swelling and discoloration		
	C. Check for pain and tenderness		
2. CARE FOR FRACTURE OF	A. Check for distal pulse		
WRIST AND FOREARM AREA	B. Check for sensation		
	C. Immobilization of fracture to wrist_and		
	forearm area using a rigid splint		
3. IMMOBILIZING A FRACTURE	A. Selection of appropriate rigid splint of		
TO WRIST AND FOREARM	proper length		
AREA			
	B. Apply appropriate padding to rigid splint		
	C. Place appropriate roller bandage in patients		
	hand to ensure the position of function		
	D. Properly apply splint with appropriate wrap		
	E. Apply wrap distal to proximal		
	F. Check for distal pulse		
	G. Check for nerve impairment		
4. IMMOBILIZING FRACTURE	A. Place sling over patients chest and under		
USING SLING	arm		
	B. MET 2 should hold or stabilize arm		
	C. Triangle should extend behind elbow on		
	injured side		
	D. Pull sling around neck until hand is		
	elevated and tie on uninjured side		
	E. Secure excess material at patients elbow		
	F. Patients finger tips should be exposed		
	G. Check for pulse and sensation		
5. SECURING SLING WITH	A. Use triangle cravat		
SWATHE			
	B. Swathe is tied around chest and injured arm		
	C. Reassess distal pulse		
	D. Reassess sensation		
	E. Treat for shock		

SPLINTING OF UPPER EXTREMITY (FRACTURES AND DISLOCATIONS)

IMMOBILIZATION OF FRACTURED RADIUS OR ULNA (USING AIR SPLINT)

RECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
1. IDENTIFICATION OF	A. Obvious signs of deformity (Often absent)		
FRACTURED RADIUS OR			
ULNA			
	B. Obvious signs of swelling and discoloration		
	C. Conscious patient check for pain and		

	tenderness	
2. CARE FOR FRACTURED	A. Check for distal pulse	
RADIUS OR ULNA		
	B. Check for sensation	
	C. Immobilize fracture with air-inflated splint	
3. IMMOBILIZE FRACTURE	A. Grasp arm with one hand above the injury	
	site with other hand just below injury site	
	B. Maintain support	
	C. Properly apply splint	
	D. Splint should be relatively free of wrinkles	
	E. Check for distal pulse prior to inflation	
	F. Inflate splint to point that slight dent can be	
	made	
	G. Check for distal pulse after inflation	
4. MONITOR AIR-INFLATED	A. Check for leaks	
SPLINT		
	B. Periodically check for increase or decrease	
	in pressure	
	C. Monitor pressure in splint with finger tip	
	D. Make certain desired pressure is	
	maintained	

TWO MAN DIRECT GROUND LIFT

RECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
1. POSITION FOR LIFT	A. Both rescuers line up on the same side of		
	the victim.		
	B. MET 1 at victim's head, and MET 2 at		
	victims legs and feet		
2. BOTH METS KNEEL ON	Rescuers place the same knee to the ground		
KNEES	(both on right knee or left knee)		
3. POSITIONING OF PATIENT	MET 1 places victim's arms across victim's		
	chest		

4. MET 1 ASSUMES PROPER POSITIONING	MET 1 cradles victim's head by placing one arm under the victim's neck and shoulder, and the other arm under victim's lower back	
5. MET 2 ASSUMES PROPER	MET 2 places one arm under the victim's	ı
POSITIONING	knees and one arm above the buttocks	
6. LIFTING THE PATIENT	On command from MET 1, both lift victim	ì
	to their knees and roll victim in toward	1
	their chest	
7. RESCUERS STAND WITH	On command from MET 1, both stand with	1
PATIENT	victim using proper body mechanics	
8. RESCUERS LOWER THE	MET 1 and MET 2 reverse procedure to	
PATIENT	lower patient as in steps 6 and 7	

ADULT 1-RESCUER CPR

RECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
1. ESTABLISH	A. Tap or gently shake shoulders		
UNRESPONSIVENESS	B. Shouts "are you okay?"		
	C. Determines unresponsiveness without		
	compromising possible c-spine injury		
	D. Call 911		
2. OPEN AIRWAY	A. Correctly executes head tilt-chin lift maneuver		
	B. Correctly executes jaw thrust maneuver, if possible spinal injuries exist		
3. CHECK FOR BREATHING	A. Look, listen and feel for air exchange for 3 - 5		
	seconds		
	B. Maintains open airway		
4. VENTILATE PATIENT	A. Seal mouth and nose with mask		

(USING ONE WAY VALVE MASK)	B. Ventilate patient: give 2 slow breaths (2 seconds per breath) – allow for chest deflation after each breath	
5. CHECK FOR CAROTID PULSE (SIGNS OF CIRCULATION)	A. Locate Adam's Apple with two fingers (index and middle fingers)	
	B. Slide fingers into grove, on same side of neck as rescuer, between windpipe and neck muscles	
	C. Feel for pulse for 5 – 10 seconds Note: If pulse is found, provide rescue breathing (1 breath every 5 seconds, about 10 to 12 breaths per minute). If no pulse is found, prepare for chest compressions	
6. POSITION FOR COMPRESSIONS	A. Kneel by victim's shoulders	
	B. Locate breastbone (sternum)	
	C. Place heel of one hand on lower half of	
	breastbone (between the nipples) and the other	
	on top of the first hand.	
	D. Keep shoulders over breastbone, with arms	
	straight	
7. DELIVER CHEST COMPRESSIONS	A. Deliver 15 smooth chest compressions (one	
	cycle) while maintaining proper body	
	positioning B. Little or no weight left on chest during upstroke	
	but contact is maintained	
8. VENTILATE PATIENT BETWEEN	A. Seal mouth and nose with mask	
COMPRESSION CYCLES	B. Ventilate patient: give 2 slow breaths (2	
(USING ONE-WAY VALVE MASK)	seconds per breath) – allow for chest	
	deflation after each breath	
9. CONTINUE CPR	A. Complete 4 cycles of 15 compressions and	
	2 breaths after each cycle	
	B. Check for presence of pulse after one	
	minute and then every few minutes	
	thereafter	

COMPLETE AIRWAY OBSTRUCTION (CONSCIOUS PATIENT)

RECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
1. INITIAL ASSESSMENT	A. Rescuer determines presence of complete airway obstruction		
	B. Ask "Are you choking?" or "Can you		
	speak?"		
2. DELIVERS ABDOMINAL THRUST	A. Positions self behind patient		
	B. Support victim with rescuers body		
	C. Wrap's arms around victim's waist		
	D. Places one fist with thumb side toward and		
	against the victim's abdomen, slightly		
	above the navel and well below the		
	xiphoid		
	E. Grasp fist with other hand		
	F. Delivers inward and upward thrust		
	(5 thrusts recommended)		
	G. Each thrust is a separate distinct movement		
	H. Thrusts are continued until foreign body is		
	expelled or becomes unconscious		
	If patient is pregnant or obese – deliver		
	chest thrusts.		
3. DELIVER CHEST THRUSTS	A. Positions self behind patient		
	B. Supports patient under armpits		
	C. Encircles victim's chest with their arms		
	while maintaining support		
	D. Places one fist with thumb side towards		
	and on the middle of the breastbone		
	E. Avoids pressure on the xiphoid and		
	margins of the rib cage		
	F. Grasp fist with other hand		
	G. Perform backward thrust until foreign		
	body is expelled or becomes unconscious		
	H. Administering thrust with the intention of		
	relieving the obstruction (5 thrusts)		
4. CONTINUE THRUSTS	Repeat thrusts until the object is expelled		
	(obstruction relieved) or the victim		
	becomes unresponsive		
	If victim becomes unresponsive (loses		
	consciousness) – continue with Skill		
	Number 17		

AIRWAY OBSTRUCTION (VICTIM LOSES CONSCIOUSNESS)

RECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
1. POSITIONING OF VICTIM	A. Rescuer places victim in supine position		
AFTER LOSS OF	taking into consideration possible c-spine		
CONSCIOUSNESS	injury		
	B. Call 911		
2. PERFORMS FINGER SWEEPS	A. Opens mouth, grasping tongue and lower		
	jaw with thumb and fingers (tongue-jaw		
	lift)		
	B. Perform a finger sweep to remove the		
	foreign object		
	C. Grasp and remove foreign object if		
2. OPENIG A IDAYAN	dislodged		
3. OPENS AIRWAY	Correctly executes head tilt - chin lift or		
A A TENENADER MENTELL A TROOP	jaw thrust maneuver		
4. ATTEMPTS VENTILATION	A. Correctly makes effort to administer 1 or 2		
NOTE: USE OF ONE WAY	rescue breaths		
VALVE MASK	B. If successful, check pulse		
VALVE MASK	C. If unsuccessful, reposition victim's head (using head tilt-chin lift or jaw thrust		
	maneuver) and attempt to administer one		
	or two more rescue breaths. If still		
	unsuccessful		
5. DELIVERS ABDOMINAL	A. Kneel astride the victims thighs		
THRUST			
	B. Places heel of one hand against the victims		
	abdomen, in mid-line slightly above the		
	navel and well below the tip of the		
	sternum (xiphoid)		
	C. Places second hand on top of first		
	D. Delivers five quick upward thrusts		
6. Repeat sequence of Functions 2. th	ru 5. until rescue breathing is successful and		
continue the steps of CPR as neede	d		

AIRWAY OBSTRUCTION (UNCONSCIOUS VICTIM - UNWITNESSED)

1. ESTABLISH UNRESPONSIVENESS A. Tap or gently shake shoulder B. Shouts "are you okay?" C. Determines unresponsiveness without compromising possible c-spine injury D. Call 911 A. Correctly executes head tilt-chin lift maneuver B. Correctly executes jaw thrust maneuver, if possible spinal injuries exist A. Look, listen and feel for air exchange for 3-5 seconds B. Maintains open airway 4. ATTEMPTS VENTILATION NOTE: USE OF ONE WAY VALVE MASK OCITECTLY makes effort to administer 1 or 2 rescue breaths B. If successful, check pulse C. If unsuccessful, reposition victim's head (using head tilt-chin lift or jaw thrust maneuver) and attempt to administer one or two more rescue breaths. If still unsuccessful A. Kneel astride the victims thighs B. Places heel of one hand against the victims abdomen, in mid-line slightly above the navel and well below the tip of the sternum (xiphoid) C. Places second hand on top of first D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and continue the steps of CPR as needed	RECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
C. Determines unresponsiveness without compromising possible c-spine injury D. Call 911 A. Correctly executes head tilt-chin lift maneuver B. Correctly executes jaw thrust maneuver, if possible spinal injuries exist A. Look, listen and feel for air exchange for 3-5 seconds B. Maintains open airway 4. ATTEMPTS VENTILATION NOTE: USE OF ONE WAY VALVE MASK B. If successful, check pulse C. If unsuccessful, reposition victim's head (using head tilt-chin lift or jaw thrust maneuver) and attempt to administer one or two more rescue breaths. If still unsuccessful 5. DELIVERS ABDOMINAL A. Kneel astride the victims thighs B. Places heel of one hand against the victims abdomen, in mid-line slightly above the navel and well below the tip of the sternum (xiphoid) C. Places second hand on top of first D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and	1. ESTABLISH	A. Tap or gently shake shoulder		
2. OPEN AIRWAY 2. OPEN AIRWAY A. Correctly executes head tilt-chin lift maneuver B. Correctly executes jaw thrust maneuver, if possible spinal injuries exist A. Look, listen and feel for air exchange for 3-5 seconds B. Maintains open airway A. Correctly makes effort to administer 1 or 2 rescue breaths NOTE: USE OF ONE WAY VALVE MASK B. If successful, check pulse C. If unsuccessful, reposition victim's head (using head tilt-chin lift or jaw thrust maneuver) and attempt to administer one or two more rescue breaths. If still unsuccessful 5. DELIVERS ABDOMINAL A. Kneel astride the victims thighs B. Places heel of one hand against the victims abdomen, in mid-line slightly above the navel and well below the tip of the sternum (xiphoid) C. Places second hand on top of first D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and	UNRESPONSIVENESS	B. Shouts "are you okay?"		
2. OPEN AIRWAY D. Call 911 A. Correctly executes head tilt-chin lift maneuver B. Correctly executes jaw thrust maneuver, if possible spinal injuries exist A. Look, listen and feel for air exchange for 3-5 seconds B. Maintains open airway A. Correctly makes effort to administer 1 or 2 rescue breaths NOTE: USE OF ONE WAY VALVE MASK B. If successful, check pulse C. If unsuccessful, reposition victim's head (using head tilt-chin lift or jaw thrust maneuver) and attempt to administer one or two more rescue breaths. If still unsuccessful A. Kneel astride the victims thighs B. Places heel of one hand against the victims abdomen, in mid-line slightly above the navel and well below the tip of the sternum (xiphoid) C. Places second hand on top of first D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and				
2. OPEN AIRWAY A. Correctly executes head tilt-chin lift maneuver B. Correctly executes jaw thrust maneuver, if possible spinal injuries exist A. Look, listen and feel for air exchange for 3-5 seconds B. Maintains open airway 4. ATTEMPTS VENTILATION NOTE: USE OF ONE WAY VALVE MASK B. If successful, check pulse C. If unsuccessful, reposition victim's head (using head tilt-chin lift or jaw thrust maneuver) and attempt to administer one or two more rescue breaths. If still unsuccessful 5. DELIVERS ABDOMINAL A. Kneel astride the victims thighs B. Places heel of one hand against the victims abdomen, in mid-line slightly above the navel and well below the tip of the sternum (xiphoid) C. Places second hand on top of first D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and				
maneuver B. Correctly executes jaw thrust maneuver, if possible spinal injuries exist A. Look, listen and feel for air exchange for 3-5 seconds B. Maintains open airway 4. ATTEMPTS VENTILATION NOTE: USE OF ONE WAY VALVE MASK C. If unsuccessful, check pulse C. If unsuccessful, reposition victim's head (using head tilt-chin lift or jaw thrust maneuver) and attempt to administer one or two more rescue breaths. If still unsuccessful 5. DELIVERS ABDOMINAL A. Kneel astride the victims thighs B. Places heel of one hand against the victims abdomen, in mid-line slightly above the navel and well below the tip of the sternum (xiphoid) C. Places second hand on top of first D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and		D. Call 911		
B. Correctly executes jaw thrust maneuver, if possible spinal injuries exist A. Look, listen and feel for air exchange for 3-5 seconds B. Maintains open airway 4. ATTEMPTS VENTILATION NOTE: USE OF ONE WAY VALVE MASK B. If successful, check pulse C. If unsuccessful, reposition victim's head (using head tilt-chin lift or jaw thrust maneuver) and attempt to administer one or two more rescue breaths. If still unsuccessful 5. DELIVERS ABDOMINAL A. Kneel astride the victims thighs B. Places heel of one hand against the victims abdomen, in mid-line slightly above the navel and well below the tip of the sternum (xiphoid) C. Places second hand on top of first D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and	2. OPEN AIRWAY	A. Correctly executes head tilt-chin lift		
possible spinal injuries exist A. Look, listen and feel for air exchange for 3-5 seconds B. Maintains open airway 4. ATTEMPTS VENTILATION NOTE: USE OF ONE WAY VALVE MASK C. If unsuccessful, check pulse C. If unsuccessful, reposition victim's head (using head tilt-chin lift or jaw thrust maneuver) and attempt to administer one or two more rescue breaths. If still unsuccessful 5. DELIVERS ABDOMINAL A. Kneel astride the victims thighs B. Places heel of one hand against the victims abdomen, in mid-line slightly above the navel and well below the tip of the sternum (xiphoid) C. Places second hand on top of first D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and				
3. CHECK FOR BREATHING A. Look, listen and feel for air exchange for 3-5 seconds B. Maintains open airway A. Correctly makes effort to administer 1 or 2 rescue breaths NOTE: USE OF ONE WAY VALVE MASK B. If successful, check pulse C. If unsuccessful, reposition victim's head (using head tilt-chin lift or jaw thrust maneuver) and attempt to administer one or two more rescue breaths. If still unsuccessful 5. DELIVERS ABDOMINAL A. Kneel astride the victims thighs B. Places heel of one hand against the victims abdomen, in mid-line slightly above the navel and well below the tip of the sternum (xiphoid) C. Places second hand on top of first D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and				
4. ATTEMPTS VENTILATION NOTE: USE OF ONE WAY VALVE MASK B. Maintains open airway A. Correctly makes effort to administer 1 or 2 rescue breaths B. If successful, check pulse C. If unsuccessful, reposition victim's head (using head tilt-chin lift or jaw thrust maneuver) and attempt to administer one or two more rescue breaths. If still unsuccessful 5. DELIVERS ABDOMINAL A. Kneel astride the victims thighs B. Places heel of one hand against the victims abdomen, in mid-line slightly above the navel and well below the tip of the sternum (xiphoid) C. Places second hand on top of first D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and				
4. ATTEMPTS VENTILATION NOTE: USE OF ONE WAY VALVE MASK B. If successful, check pulse C. If unsuccessful, reposition victim's head (using head tilt-chin lift or jaw thrust maneuver) and attempt to administer one or two more rescue breaths. If still unsuccessful 5. DELIVERS ABDOMINAL A. Kneel astride the victims thighs B. Places heel of one hand against the victims abdomen, in mid-line slightly above the navel and well below the tip of the sternum (xiphoid) C. Places second hand on top of first D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and	3. CHECK FOR BREATHING			
4. ATTEMPTS VENTILATION NOTE: USE OF ONE WAY VALVE MASK C. If unsuccessful, check pulse C. If unsuccessful, reposition victim's head (using head tilt-chin lift or jaw thrust maneuver) and attempt to administer one or two more rescue breaths. If still unsuccessful 5. DELIVERS ABDOMINAL A. Kneel astride the victims thighs B. Places heel of one hand against the victims abdomen, in mid-line slightly above the navel and well below the tip of the sternum (xiphoid) C. Places second hand on top of first D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and				ļ
NOTE: USE OF ONE WAY VALVE MASK C. If unsuccessful, reposition victim's head (using head tilt-chin lift or jaw thrust maneuver) and attempt to administer one or two more rescue breaths. If still unsuccessful 5. DELIVERS ABDOMINAL A. Kneel astride the victims thighs B. Places heel of one hand against the victims abdomen, in mid-line slightly above the navel and well below the tip of the sternum (xiphoid) C. Places second hand on top of first D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and				ļ
NOTE: USE OF ONE WAY VALVE MASK C. If unsuccessful, reposition victim's head (using head tilt-chin lift or jaw thrust maneuver) and attempt to administer one or two more rescue breaths. If still unsuccessful 5. DELIVERS ABDOMINAL A. Kneel astride the victims thighs B. Places heel of one hand against the victims abdomen, in mid-line slightly above the navel and well below the tip of the sternum (xiphoid) C. Places second hand on top of first D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and	4. ATTEMPTS VENTILATION			
VALVE MASK C. If unsuccessful, reposition victim's head				-
(using head tilt-chin lift or jaw thrust maneuver) and attempt to administer one or two more rescue breaths. If still unsuccessful 5. DELIVERS ABDOMINAL A. Kneel astride the victims thighs B. Places heel of one hand against the victims abdomen, in mid-line slightly above the navel and well below the tip of the sternum (xiphoid) C. Places second hand on top of first D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and		*		-
maneuver) and attempt to administer one or two more rescue breaths. If still unsuccessful 5. DELIVERS ABDOMINAL A. Kneel astride the victims thighs B. Places heel of one hand against the victims abdomen, in mid-line slightly above the navel and well below the tip of the sternum (xiphoid) C. Places second hand on top of first D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and	VALVE MASK			
or two more rescue breaths. If still unsuccessful 5. DELIVERS ABDOMINAL A. Kneel astride the victims thighs B. Places heel of one hand against the victims abdomen, in mid-line slightly above the navel and well below the tip of the sternum (xiphoid) C. Places second hand on top of first D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and				
5. DELIVERS ABDOMINAL A. Kneel astride the victims thighs B. Places heel of one hand against the victims abdomen, in mid-line slightly above the navel and well below the tip of the sternum (xiphoid) C. Places second hand on top of first D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and				
5. DELIVERS ABDOMINAL A. Kneel astride the victims thighs B. Places heel of one hand against the victims abdomen, in mid-line slightly above the navel and well below the tip of the sternum (xiphoid) C. Places second hand on top of first D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and				
B. Places heel of one hand against the victims abdomen, in mid-line slightly above the navel and well below the tip of the sternum (xiphoid) C. Places second hand on top of first D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and	5 DELIVERS ARROMINAL			
abdomen, in mid-line slightly above the navel and well below the tip of the sternum (xiphoid) C. Places second hand on top of first D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and	3. DELIVERS ABDOMINAL			
navel and well below the tip of the sternum (xiphoid) C. Places second hand on top of first D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and				
sternum (xiphoid) C. Places second hand on top of first D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and				
C. Places second hand on top of first D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and		=		
D. Delivers five quick upward thrusts 6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and				+
6. Repeat sequence of Functions 2. thru 5. until rescue breathing is successful and				+

ADULT 2-RESCUER CPR

RECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
1. ESTABLISH UNRESPONSIVENESS	A. Tap or gently shake shoulder		
	B. Shouts "are you okay?"		
	C. Determines unresponsiveness without		
	compromising possible c-spine injury		
	D. Call 911		
Note: Rescuer Number 2 (rescuer remaini			
2. OPEN AIRWAY	A. Correctly executes head tilt – chin lift		
(RESCUER NUMBER 2)	maneuver		
	B. Correctly executes jaw thrust maneuver, if		
	possible spinal injuries exist		
3. CHECK FOR BREATHING	A. Look, listen and feel for air exchange for		
(RESCUER NUMBER 2)	3-5 seconds		
	B. Maintains open airway		
4. VENTILATE PATIENT	A. Seal mouth and nose with mask		
(USING ONE WAY VALVE MASK)	B. Ventilate patient: give 2 slow breaths (2		
·	seconds per breath) – allow for chest deflation		
(RESCUER NUMBER 2)	after each breath		
5. CHECK FOR CAROTID PULSE	A. Locate Adam's Apple with two fingers (index		Ì
(SIGNS OF CIRCULATION)	and middle fingers)		
(RESCUER NUMBER 2)	B. Slide fingers into grove, on same side of neck		
,	as rescuer, between windpipe and neck		
	muscles		
	C. Feel for pulse for 5 – 10 seconds Note: If		
	pulse is found, provide rescue breathing (1		
	breath every 5 seconds, about 10 to 12 breaths		
	per minute). If no pulse if found, prepare for		
	chest compressions		
Note: Both rescuers begin working in unis			
6. POSITION FOR COMPRESSIONS	A. Kneel by victim's shoulders		
(RESCUER NUMBER 1)	B. Locate breastbone (sternum)		
	C. Place heel of one hand on lower half of		
	breastbone (between the nipples) and the other		
	on top of the first hand		
	D. Keep shoulders over breastbone, with arms		
	straight		
7. DELIVER CHEST	A. Deliver 15 smooth chest compressions (one		
COMPRESSION	cycle) while maintaining proper body		
(RESCUER NUMBER 1)	positioning		
	B. Little or no weight left on chest during upstroke		
	but contact is maintained		
8. VENTILATE PATIENT BETWEEN	A. Seal mouth and nose with mask		
COMPRESSION CYCLES	B. Ventilate patient: give 2 slow breaths (2		
(USING ONE-WAY VALVE MASK)	seconds per breath) – allow for chest deflation		
(RESCUER NUMBER 2)	after each breath		
9. CONTINUE CPR	Complete 4 cycles of 15 compressions and 2		
(BOTH RESCUERS)	breaths after each compression cycle		
10. CHECK PULSE AND	Check for presence of pulse after one minute		
BREATHING	and then every few minutes thereafter (check		
(RESCUER 2)	for breathing simultaneously)		

ESTABLISHING AIRWAY, SUSPECTED CERVICAL INJURY

RECOMMENDED FUNCTION	CRITICAL SKILL	Y	N
1. STABILIZE HEAD	Restrains victim's head and neck to		
	avoid voluntary or involuntary		
	movement causing flexion,		
	hyperextension rotation or lateral		
	movement of the victims neck		
2. OPEN THE AIRWAY	A. Use modified jaw thrust maneuver		
	without causing hyperextension of		
	victims neck		
	B. There should be no movement of the		
	victims head or neck that would		
	compromise the suspected cervical		
	injury		
3. CHECKS FOR BREATHING	A. Rescuer takes a full 5 seconds to		
	check		
	for breathing		
	B. States, victim is breathing		
4. MAINTAIN OPEN AIRWAY	Do not compromise suspected cervical		
	injury		